

UBVRI photometric sequences for symbiotic stars

Ulisse Munari

*Osservatorio Astronomico di Padova – INAF, Sede di Asiago, I-36012
 Asiago (VI), Italy*

Arne Henden

*Universities Space Research Association/U.S. Naval Observatory
 Flagstaff Station, P.O. Box 1149, Flagstaff AZ 86002-1149, USA*

Abstract. Deep, wide-range and accurate UBVRI photometric sequences have been established around more than 80 symbiotic stars, to assist current photometry as well as measurement of old photographic plates. Sequences for 40 symbiotic stars have already been published; the observations for the others have already been secured.

1. Introduction

Availability of suitable UBVRI photometric sequences around symbiotic stars is an essential step in promoting a large-scale documentation effort of present time multi-band lightcurves as well as reconstruction of the historical behavior.

To serve for measuring archival plates, the sequence stars have to be well isolated from neighbors to avoid blending on short focal length patrol plates, have to be grouped close to the symbiotic variable so as to fall within the same eyepiece field of view, and must extend over a wide range in brightness to easily cover both faint states in quiescence (like eclipses) as well as bright outbursts.

To assist present time CCD photometry (carried out mostly by amateurs with telescopes of 1-2 m focal lengths), the sequences must also cover a wide range in colors (to allow simultaneous derivation of transformation coefficients), be compact enough to easily fit onto a small CCD ($\lesssim 5$ arcmin in diameter) and be accurately placed on the UBVRI system.

The UBVRI sequences on the Johnson-Cousins system that we have calibrated around more than 80 symbiotic stars follow the above prescriptions and are discussed in this paper. Their use on Asiago archive plates is described by Jurdana-Sepic and Munari (2002, and references therein).

2. The sequences

Figure 1 and Table 1 show a typical UBVRI photometric sequence among the 82 we have calibrated. The increasing errors at the brightest magnitudes are due to saturation effects, and those at the faintest magnitudes are dominated by the photon statistics.




Figure 1. Identification chart for the UBVRI sequence in Table 1.

The symbiotic stars for which we have already obtained the multi-epoch UBVRI observations suitable for the calibration of the photometric sequences are listed in Table 2. Of the 82 listed objects, 40 have their sequences published by Henden and Munari (2000, 2001); the remaining 42 will follow shortly. The program stars have been mainly selected from the catalogues of symbiotic stars of Allen (1984) and Belczyński et al. (2000)

All observations were made with the 1.0 m Ritchey-Chrétien telescope of the U.S. Naval Observatory, Flagstaff Station. A Tektronix/SITe 1024×1024 thinned, backside-illuminated CCD was used, along with proper Johnson UBV and Kron-Cousins RI filters. The telescope scale is 0.67 arcsec/pix and typical seeing was ~ 2 arcsec. The color transformation equations for this instrumental combination are particularly good, with slopes deviating from unity by only a few hundredths of a magnitude (see Eq.(1)-(5) in Henden and Munari 2000). Only data collected under photometric conditions (transformation errors under

	N	B	V	B-V	U-B	V-R	R-I
a	3	12.896 $\pm$ 0.012	11.632 $\pm$ 0.023	1.264 $\pm$ 0.035	1.092 $\pm$ 0.045		
b	3	13.510 $\pm$ 0.003	12.782 $\pm$ 0.005	0.728 $\pm$ 0.003	0.256 $\pm$ 0.016	0.402 $\pm$ 0.001	0.394 $\pm$ 0.002
c	3	13.838 $\pm$ 0.002	13.289 $\pm$ 0.005	0.549 $\pm$ 0.001	-0.009 $\pm$ 0.016	0.318 $\pm$ 0.000	0.334 $\pm$ 0.009
d	3	14.799 $\pm$ 0.004	13.092 $\pm$ 0.005	1.707 $\pm$ 0.001	1.698 $\pm$ 0.008	0.950 $\pm$ 0.006	0.891 $\pm$ 0.007
e	3	15.426 $\pm$ 0.005	14.486 $\pm$ 0.006	0.940 $\pm$ 0.005	0.705 $\pm$ 0.003	0.533 $\pm$ 0.011	0.484 $\pm$ 0.005
f	3	16.025 $\pm$ 0.004	15.108 $\pm$ 0.006	0.917 $\pm$ 0.004	0.263 $\pm$ 0.007	0.555 $\pm$ 0.007	0.563 $\pm$ 0.003
g	3	16.546 $\pm$ 0.000	15.528 $\pm$ 0.000	1.018 $\pm$ 0.006	0.457 $\pm$ 0.010	0.608 $\pm$ 0.002	0.615 $\pm$ 0.010
h	3	16.900 $\pm$ 0.004	15.115 $\pm$ 0.000	1.785 $\pm$ 0.006	1.858 $\pm$ 0.009	1.016 $\pm$ 0.009	0.964 $\pm$ 0.008
l	3	17.468 $\pm$ 0.002	16.335 $\pm$ 0.006	1.133 $\pm$ 0.009	1.018 $\pm$ 0.021	0.667 $\pm$ 0.004	0.597 $\pm$ 0.003
m	3	18.108 $\pm$ 0.006	16.997 $\pm$ 0.009	1.111 $\pm$ 0.005	0.629 $\pm$ 0.040	0.640 $\pm$ 0.013	0.620 $\pm$ 0.015
n	3	19.057 $\pm$ 0.010	17.884 $\pm$ 0.009	1.173 $\pm$ 0.020	0.527 $\pm$ 0.074	0.726 $\pm$ 0.012	0.660 $\pm$ 0.053
z	3	20.072 $\pm$ 0.017	18.509 $\pm$ 0.026	1.563 $\pm$ 0.043		1.016 $\pm$ 0.005	1.050 $\pm$ 0.071

Table 1. The UBVRI comparison sequence plotted in Figure 1. N is the number of observations on separate nights used to check for variability. The quoted errors are formal ones.

Z And	CI Cyg	RS Oph	V4368 Sgr	Hen 3-1341	Wray 1470
EG And	V407 Cyg	V1261 Ori	IV Vir	Hen 3-1342	Wray 157
CM Aql	V1016 Cyg	AG Peg	PU Vul	Hen 3-1591	
V352 Aql	V1329 Cyg	AX Per	V335 Vul	K 3-9	
V1413 Aql	ER Del	V471 Per	ALS 1	M 1-21	
R Aqr	LT Del	DQ Ser	ALS 2	MWC 960	
UV Aur	AG Dra	DT Ser	AS 201	Mac 1-17	
TX CVn	GH Gem	FG Ser	AS 210	NSV 11776	
BD Cam	NQ Gem	RT Ser	AS 289	Pe 2-16	
CI Cam	YY Her	UU Ser	AS 323	Pt 1	
LL Cas	V443 Her	HM Sge	AS 327	StHa 149	
V627 Cas	V503 Her	QW Sge	Ap 3-1	StHa 164	
ZZ CMi	V934 Her	FN Sgr	UKS Ce1	StHa 169	
T CrB	RW Hya	V919 Sgr	Draco C-1	StHa 180	
BF Cyg	BX Mon	V4018 Sgr	Hen 2-442	StHa 190	
CH Cyg	V694 Mon	V4074 Sgr	Hen 2-468	StHa 32	

Table 2. List of symbiotic stars for which we have obtained a UBVRI photometric sequence. Forty of them have been published in Henden and Munari (2000, 2001); for the remaining 42, the observations have been completed and publication will follow.

0.02 mag) have been used, and the calibration has been made against the Landolt equatorial fields (Landolt 1983, 1992).

Each photometric sequence is composed of 10 to 15 stars, which have been selected and ordered on the basis of their B band magnitude. The latter is reproducible by most filter-equipped CCD cameras, it is the closest one to the m_{pg} band of historical observations, and the B band is particularly well suited to investigate the variability of symbiotic stars. Stars of varying color are included in the sequence to allow simultaneous determination of transformation coefficients. The comparison sequences are tighter around the usual brightness of each symbiotic star and become looser away from it.

The stars included in the comparison sequences have been checked on at least three nights for variability. We cannot obviously rule out beyond doubt that some of them are indeed variable, but the agreement at a few milli-mag level of their magnitude measured on different nights over an interval of months gives some confidence in their use.

References

- Allen, D.A. 1984, Proc. A.S.A., 5, 369
Belczyński K., Mikolajewska J., Munari U., Ivison R.J. & Friedjung M. 2000, A&AS, 146, 407
Jurdana-Sepic, R., Munari, U., 2002, this volume
Henden A., Munari U. 2000, A&AS143, 343
Henden A., Munari U. 2000, A&A372, 145
Landolt, A. U., 1983, AJ88,439.
Landolt, A. U., 1992, AJ104,340.